



SURGICAL SITE INFECTIONS IN GENERAL SURGICAL WARDS AT TERTIARY CARE HOSPITAL CENTRAL REGION OF INDIA: ANALYTICAL CROSS-SECTIONAL STUDY

**Dr Sonia
Shrivastava***

Assistant Professor Department Of Microbiology L.M.N.C Bhopal *Corresponding Author

ABSTRACT

Background: Surgical site infections are important cause of morbidity and mortality in admitted patients world over. **Objectives:** To determine the pattern of surgical site infections in General Surgical ward of a tertiary care hospital. **Study type, settings and duration:** Analytical cross-sectional study conducted at Department of General Surgery, LAXMI NARAYAN MEDICAL COLLEGE & RESEARCH CENTRE, and BHOPAL me for two years from January 2017 to December 2021. **Subjects and Methods:** All cases were admitted in surgical ward with various surgical problems either as elective or emergency cases who developed wound infection later were included in the study. Cases of wound infection operated elsewhere, diabetic foot, and abscesses were excluded. Data collected included age, gender, primary diagnosis, mode of admission, comorbid factors, type and duration of surgery, expertise of the surgeon, use of antibiotics and hospital stay. After operation, wound was examined for evidence of infection from third post-op day onward. Any discharge was submitted for bacteriological examination. The wounds were followed till healed. **Results:** A total of 478.25 patients underwent surgery, including 245.75 cases (51.5 %) operated as elective and 233(48.5 %) as emergency. Postoperative wound infections occurred in 41.5 cases giving an overall incidence of infection in 8.6% cases. Infection rate in elective cases was lower (4.6%) than that in the emergency (12.7%) cases. Sixty one patients (37%) developed minor infection or stitch abscess, 26(63%) has frank suppuration requiring opening and drainage of wound; while 1.2(3%) cases developed deep seated infection of intra-abdominal spaces. E. coli was the commonest bacteria for wound infection (39%). **Conclusion:** Post operative wound infection rate was 8.6%. The infection was significantly higher in cases who underwent emergency surgery and E. coli was the commonest pathogen to cause infection.

KEYWORDS

Surgical site infections, surgical wound infections, bacterial infections, antibiotic prophylaxis.

Corresponding Author Dr Sonia Shrivastava Assistant Professor Department Of Microbiology L.M.N.C Bhopal *Corresponding Author

Introduction

Infections which develop more than 48 hours after admission are hospital acquired or nosocomial infections¹

They are the sixth leading cause of death in USA, accounting for 150,000 deaths per year, and their incidence varies from 5-10%². Nosocomial infections are an important cause of preventable morbidity and mortality that prolong hospital stay by an average of eight days³

while the treatment cost and work load on health care facilities is increased many folds. Among surgical patients, surgical site infections (SSIs) are one of the most c SSIs cdebilitate patients and dramatically increase health care costs⁴

They are a leading cause of readmission, may lead to complications like delayed wound healing and revision surgery⁷ and with longer hospital stay can render patients susceptible to infections from the hospital environments⁸.

The CDC-definitions for surveillance of surgical site infections take into account 3 classes of wound infections: superficial, deep incisional SSI, and organ/space SSI⁹. Since ancient times wound infections have markedly increased the sufferings of postoperative cases: and despite being largely preventable, they remain a major source of morbidity¹⁰. In order to minimize postoperative surgical wound infection, it is important to create a safe environment by controlling four main sources of infection i.e. personnel, equipment, the environment, and patient`s risk factors¹¹.

Knowledge of specific risk factors for SSI is essential to create a specific SSI risk stratification index¹², and to develop strategies to confine infection rate. The best approach is the prevention as it is simpler, cheaper and more rewarding for the patients; and at least one third of them are preventable by simple measures¹³.

Thus, every hospital needs to organize its infection control program. Failure to implement infection control policies and lack of awareness

are the factors contributing to hospital infections and disease outbreaks. On the other hand, studies provide evidence of a significant decreasing trend in the SSI rates following the infection control interventions¹⁴.

In our hospitals, there are high rates of nosocomial infections with little efforts to control them. The present study was conducted on patients operated for various surgical problems in a tertiary care major public sector of Bhopal to evaluate the frequency of surgical site infections.

Materials and Methods

This analytical cross-sectional study was conducted in the department of general surgery, LAXMI NARAYAN MEDICAL COLLEGE & RESEARCH CENTRE, BHOPAL over two years from January 2010 to December 2011. All adult cases admitted in surgical ward either as elective cases or as emergency were included in the study. Cases having wound infection, operated elsewhere, those with diabetic foot disease and those operated for incision and drainage of abscess were excluded. All the cases operated during the study period were observed for development of wound infection. Patients undergoing multiple admissions or operations for complications were counted once.

Consent was taken from all the patients who developed infection, for inclusion in the study. All the information collected was recorded on a specially designed proforma that included history, physical examination, any co-morbid factors and all investigations performed during their stay at the hospital. The operative details were noted including type of procedure, duration of surgery, whether operated by residents or consultants, and use of prophylactic antibiotics. The wounds were examined for infection from third postoperative day onward. Surgical site infection was identified with redness, inflammation, local heat, pain, temperature of 38° C or above, and septic discharge from incision site during 30 days after operation (according to WHO guidelines)¹⁵

The discharge was sent for culture and sensitivity. The infections were managed by standard protocol involving repeated dressings according

to the case, drainage of pus if required, and change of antibiotic in the light of culture/sensitivity reports. Pre and postoperative hospital stay was noted. Infected wounds were inspected regularly during follow up until they were healed.

The information was entered in computer and data was analyzed using SPSS version 12.0. Various frequencies and percentages were calculated; the results are displayed in tabulated or graphic forms.

Results

During the study period, a total of 527 cases were registered in surgical ward and 478.75 were operated. Surgery in 245.75 cases (51.3%) was elective or planned while 233 patients (48.7 %) were operated in emergency. Majority were males 291.75(61%). The median age was 42.7 years (range 16-82 years) and majority 354.25(74%) were below 50 years of age. Almost 58% men developed infection. Postoperative infection occurred in 41.25 cases giving an overall incidence of 8.6% infection. Of 41.25 cases that developed wound infection, 17.5 (42%) patients were aged above 50 years. Infections were least common 5(2.1%) in clean procedures and were highest among dirty cases (Table-1). Out of 245.75 cases who were operated on elective list, infection occurred in 11.5 patients (4.6%). Out of 233 cases operated in emergency, infection occurred in 29.75 patients (12.7%).

Table 1: Infection rate in different procedures.

| Type of procedure | Total cases (n=478.75) | | Infected cases (41.25) | |
|-----------------------------|--------------------------|------|------------------------|------|
| | n | % | n | % |
| Clean | 244 | 51 | 5 | 2.1 |
| Clean-contaminated | 105.25 | 22 | 4.75 | 4.5 |
| Contaminated | 43.25 | 9 | 7 | 16.2 |
| Dirty | 86.25 | 18 | 24.5 | 28.5 |
| Operative settings | | | | |
| Setting | Number of infected cases | % | | |
| Elective surgery (n=245.75) | 11.5 | 4.6 | | |
| Emergency surgery (n=233) | 29.75 | 12.7 | | |

Average duration of surgery was 109 minutes (range 35 minutes to 7 hours 20 minutes). Antibiotic prophylaxis was given in clean cases (first generation cephalosporins) and clean-contaminated cases (third generation cephalosporins). In contaminated or dirty cases, regular use of antibiotics was employed according to the case, rather than giving prophylaxis.

Average duration of hospital stay after surgery was 7.4 days (range 1-110 days), while average duration of hospital stay before surgery was 1.8 days (range 10 hours to 13 days). Although infected cases took longer in hospital, however, this aspect was not studied in detail. Of the co-morbid, anemia (haemoglobin less than 10 gm/dl) was the most common. In elective cases it was corrected before surgery but in emergency cases per operative blood transfusion was given. Other co-morbid factors in 41.25 patients are shown in Table-2.

Surgical wound infection developed in 41.25 cases. Majority 24.75(60%) showed frank suppuration that required opening and drainage of wound. Wound infection was observed within 3-5 days in 27.5 cases (67%), within 6-8 days in 10.25 cases (25%), and within 9-14 days in 3.25 cases (8%). One case presented with infection 10 months after mesh repair for incisional hernia.

Table 2: Distribution of co-morbid factors. (n=41.25)

| Factor observed | n | % |
|---|-------|------|
| Anemia (Haemoglobin less than 10 gm/dl) | 15.75 | 38 |
| Malnutrition (Loss of more than 6 Kg body weight) | 10.25 | 25 |
| Smoking (more than 10 cigarettes/day) | 6.5 | 16 |
| Diabetes mellitus | 4.75 | 11.5 |
| Hypertension and / or ischemic heart disease. | 4.75 | 11.5 |
| Obesity (more than 10% of ideal body weight) | 2 | 5.0 |
| Compensated chronic liver disease | 0.75 | 2.0 |

Discharge from the wound was submitted for bacterial culture in all cases. Twenty three cases (14%) showed mixed growth; 121(73%) showed growth of single organism; and 21(13%) had no growth. The commonest bacteria grown was E. coli in 64(39%) cases, Klebsiella spp. in 37(22%) cases, Pseudomonas aeruginosa in 25(15%) cases, and *Styphlococcus aureus* in 18(11%) cases including methicillin resistant *S. aureus* (MRSA) (Figure-1).




Figure 1: Organisms grown on culture.

Discussion

In the present study surgical site infection were seen in 8.6% cases with majority of the cases undergoing emergency surgery. The incidence of SSI varies and is 4.4% in Taiwan¹⁶, 5% in United States¹⁷, and 5.2% in Japan¹⁸, while a Brazilian study reported a much lower incidence of 1.8%¹⁹. Reports from Pakistan show a higher incidence of 6.5% from Peshawar²⁰, and 11.4% from Karachi²¹.

Several factors are responsible for causing infections which vary from patients themselves (especially contamination by alimentary tract bacteria) to other patients, hospital environment, food, hospital staff, infected surgical instruments, dressings, and even medicines and injections²². Advanced age is an important host-related risk factor²². Due to higher incidence of coexisting diseases, impaired immunological status, personal neglect, etc. However, gender is not a significant issue and same has been reported by others^{22,23}.

Type of surgery is directly related to the risk of developing wound infection. It is based on potential bacterial contamination of the tissues at the time of surgery and the level of bacterial burden. Our study also showed the association of type of surgery with infection rate and same reported by other worker²². Duration of surgery also influences wound infection and procedures that take more than two hours are associated with higher infection rates²⁴, due to longer exposure of tissues to theater environment, hypothermia²⁵, and requirement of blood transfusion all of them are potential risk factors for SSI²⁶. In a study from Lahore, wound infection rate almost double in cases that took longer than 2 hours showing direct relation to duration of surgical procedure²⁴. This was also observed in our study where most of the cases that got infected took longer than 100 minutes.

Operative settings (elective or emergency) also play a significant role in determining infection rates. Cases operated in emergency are more likely to get infected due to inadequate preparation, breach in sterilization protocol, pre-existing infection and reduced immunological status of patient. In the present study, infection rate in emergency cases was almost three times higher than in elective cases (12.7% versus 4.6%). This has also been observed by other workers. Studies from Lahore show about two and a half²⁷ to four times²⁴ higher infection rate in emergency cases.

Longer hospital stay, especially in post-operative period, is associated with substantial increase in wound infection rates²²; and the risk increases with the duration of stay. On the other hand, prolonged preoperative hospital stay also contributes to increased infection rate⁹. This may be related to bacterial colonization of patient's skin and nares with resistant hospital flora.

Expertise of surgeon is a potential factor in determining wound infection rate as reported by some authors²⁶. We did not observe any significant difference of infection rate between cases operated by seniors and juniors. Improvement in the surgical skills and techniques of resident staff and also their direct supervision not only decreases the duration of operation but also incidence of postoperative wound infection²⁴.

In our study most common bacterial growth was of E. coli, followed by *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Styphlococcus aureus* including MRSA. Similar pattern has also been reported from Hyderabad²⁵, but this is in contrast to the literature that reported *Styphlococcus aureus* as the most common organism^{20,26-28}, that is resistant to the commonly used antibiotics.

References

1. Khurshid A. Hospital-acquired infections. J Pak Inst Med Sci

- 2004;15:886-9.2. Emmerson AM, Enstone JE, Griffen H, Kelsey MC, Smyth ETM. The second national Prevalence survey of infection in hospitals. Overview of the results. *J Hosp Infect* 1996; 32: 175-90.
3. Badaruddin AM. Nosocomial infections; urgent need for structured and coherent approach to the problem in Pakistan. *Prof Med J* 2007;14:70-5.
 4. Smyth ET, Emmerson AM. Surgical site infection surveillance. *J Hosp Infect.* 2000;45:173-84.
 5. Cruse PJ, Foord R. The epidemiology of wound infection: a 10-year prospective study of 62,939 wounds. *Surg Clin N Am* 1980; 60:27-40.
 6. Olsen MA, Chu-Ongsakul S, Brandt KE, Dietz JR, Mayfield J, Fraser VJ. Hospital-associated costs due to surgical site infection after breast surgery. *Arch Surg.* 2008 Jan;143(1):53-60.
 7. Harrop JS, Styliaras JC, Ooi YC, Radcliff KE, Vaccaro AR, Wu C. Contributing factors to surgical site infections. *J Am Acad Orthop Surg.* 2012 Feb;20(2):94-101.
 8. Mehrun N, Tallat N, Irum A, Lubna H: Scope of surgical site infection (SSI) in Obstetrics and Gynecology. *J Postgrad Med Inst Oct - Dec 2005;19(4):438-41.*
 9. Lauwers S, de Smet F. Surgical site infections. *Acta Clin Belg.* 1998 Oct;53(5):303-10.
 10. Coakley BA, Divino CM. Identifying factors predictive of surgical-site infections after colectomy for fulminant ulcerative colitis. *Am Surg.* 2012 Apr;78(4):481-4.
 11. Damani NN, Mughis Uddin Ahmed. Prevention of Surgical Wound Infection. *Ann Abbasi Shaheed Hosp Karachi Med Dent Coll Jan 1999;4:131-2.*
 12. Anannamcharoen S, Vachirasrisirikul S, Boonya-Assadorn C. Incisional surgical site infection in colorectal surgery patients. *J Med Assoc Thai.* 2012 Jan;95(1):42-7.
 13. Haley RW, Culver DH, White JW. The efficacy of infection surveillance and control Programmes in preventing nosocomial infections in US hospitals. *Am J Epidemiol* 1985;121:182-205.
 14. Barchitta M, Matranga D, Quattrocchi A, Bellocchi P, Ruffino M, Basile G, Agodi A. Prevalence of surgical site infections before and after the implementation of a multimodal infection control programme. *J Antimicrob Chemother.* 2012 Mar;67(3):749-55.
 15. Spencer FC: Bacterial infections (in) Schwartz SI, Comshires G, Spencer FC, Dally GN, Fischer J, Galloway AC (eds): Principles of surgery. 7th edn. New York, McGraw-Hill, 1999:83.
 16. Chuang SC, Lee KT, Chang WT, Wang SN, Kuo KK, Chen JS, et al: Risk factors for wound infection after cholecystectomy. *J Formos Med Assoc.* 2004;103(8):607-12.
 17. William GC: Risk factors for surgical site infection. *Surg Infec* 2006;Vol.7:57-11
 18. Konishi T, Hariharay, Morikane K: Surgical site infection surveillance. *Nippon Geka Gakkai Zasshi.* 2004;105:720-5.
 19. Ercole FF, Franco LM, Macieira TG, Wenceslau LC, de Resende HI, Chianca TC. Risk of surgical site infection in patients undergoing orthopedic surgery. *Rev Lat Am Enfermagem.* 2011 Nov-Dec;19(6):1362-8.
 20. Nisa M, Naz T, Afzal I, Hassan L. Scope of surgical site infection (SSI) in obstetrics and gynecology. *J Post Med Ins* 2005;19:438-41.
 21. Jawaid M, Masood Z, Iqbal SA. Post-operative complications in a general surgical ward of a Teaching Hospital. *Pak J Med Sci* 2006;22(2):171-5.
 22. Razavi SM, Ibrahimpoor M, Kashani AS, Jafarain A: Abdominal surgical site infections: incidence and risk factors at an Iranian teaching hospital. *BMC Surg* 2005;5:2-7.
 23. Habte GE, Gedebau M, Kronvall G: Hospital acquired infections among surgical patients in Tikur Anbessa Hospital, Addis Ababa, Ethiopia. *Am J Infect Control* 1988:7-13.
 24. Akhtar S, Ali A, Farhan K, Nadeem A, Gondal KM, Majeed C. Duration of surgery: does it contribute to post-operative wound infection? *Pak J Surg* 2001;17:35-40.
 25. Asad A, Tahir SM, Memon AS, Noshad AS: Pattern of pathogens and their sensitivity isolated from superficial surgical site infections in a tertiary care hospital. *J Ayub Med Coll Abbottabad* 2009;21:80-2.
 26. Madu KA, Enweani UN, Katchy AU, Madu AJ, Aguwa EN. Implant associated surgical site infection in orthopaedics: a regional hospital experience. *Niger J Med.* 2011;20:435-40.
 27. Ghazi MA, Akram Y, Cheema MA. Sensitivity and resistance of micro-organisms in infected surgical wounds on surgical floor of Mayo Hospital. *Ann King Edward Med Coll* 2005;11:295-8.
 28. Ramos N, Skeete F, Haas JP, Hutzler L, Slover J, Phillips M, Bosco J. Surgical site infection prevention initiative - patient attitude and compliance. *Bull NYU Hosp Jt Dis.* 2011;69:312-5.